

Recovery Strategy for the Fascicled Ironweed (*Vernonia fasciculata*) in Canada

Fascicled Ironweed



2021



2 **Recommended citation:**
3

4 Environment and Climate Change Canada. 2021. Recovery Strategy for the Fascicled
5 Ironweed (*Vernonia fasciculata*) in Canada [Proposed]. *Species at Risk Act Recovery*
6 *Strategy Series*. Environment and Climate Change Canada, Ottawa. vii + 36 pp.

7
8
9
10 **Official version**

11 The official version of the recovery documents is the one published in PDF. All
12 hyperlinks were valid as of date of publication.

13
14 **Non-official version**

15 The non-official version of the recovery documents is published in HTML format and all
16 hyperlinks were valid as of date of publication.

17
18
19
20 For copies of the recovery strategy, or for additional information on species at risk,
21 including the Committee on the Status of Endangered Wildlife in Canada (COSEWIC)
22 Status Reports, residence descriptions, action plans, and other related recovery
23 documents, please visit the [Species at Risk \(SAR\) Public Registry](#)¹.

24
25
26
27 **Cover illustration:** © Candace Neufeld

28
29
30 Également disponible en français sous le titre
31 « Programme de rétablissement de la vernonie fasciculée (*Vernonia fasciculata*) au
32 Canada [Proposition] »

33
34 © Her Majesty the Queen in Right of Canada, represented by the Minister of
35 Environment and Climate Change, 2021. All rights reserved.

36 ISBN
37 Catalogue no.

38
39
40 *Content (excluding the illustrations) may be used without permission, with appropriate*
41 *credit to the source.*

¹ www.canada.ca/en/environment-climate-change/services/species-risk-public-registry.html

42 Preface

43
44 The federal, provincial, and territorial government signatories under the [Accord for the](#)
45 [Protection of Species at Risk \(1996\)](#)² agreed to establish complementary legislation and
46 programs that provide for effective protection of species at risk throughout Canada.
47 Under the *Species at Risk Act* (S.C. 2002, c.29) (SARA), the federal competent
48 ministers are responsible for the preparation of recovery strategies for listed Extirpated,
49 Endangered, and Threatened species and are required to report on progress within
50 five years after the publication of the final document on the SAR Public Registry.

51
52 The Minister of the Environment and Climate Change Canada is the competent minister
53 under SARA for the Fascicled Ironweed and has prepared this recovery strategy, as per
54 section 37 of SARA. To the extent possible, it has been prepared in cooperation with
55 the Government of Manitoba as per section 39(1) of SARA.

56
57 Success in the recovery of this species depends on the commitment and cooperation of
58 many different constituencies that will be involved in implementing the directions set out
59 in this strategy and will not be achieved by Environment and Climate Change Canada,
60 or any other jurisdiction alone. All Canadians are invited to join in supporting and
61 implementing this strategy for the benefit of the Fascicled Ironweed and Canadian
62 society as a whole.

63
64 This recovery strategy will be followed by one or more action plans that will provide
65 information on recovery measures to be taken by Environment and Climate Change
66 Canada and other jurisdictions and/or organizations involved in the conservation of the
67 species. Implementation of this strategy is subject to appropriations, priorities, and
68 budgetary constraints of the participating jurisdictions and organizations.

69
70 The recovery strategy sets the strategic direction to arrest or reverse the decline of the
71 species, including identification of critical habitat to the extent possible. It provides all
72 Canadians with information to help take action on species conservation. When critical
73 habitat is identified, either in a recovery strategy or an action plan, SARA requires that
74 critical habitat then be protected.

75
76 In the case of critical habitat identified for terrestrial species including migratory birds
77 SARA requires that critical habitat identified in a federally protected area³ be described
78 in the *Canada Gazette* within 90 days after the recovery strategy or action plan that
79 identified the critical habitat is included in the public registry. A prohibition against
80 destruction of critical habitat under ss. 58(1) will apply 90 days after the description of
81 the critical habitat is published in the *Canada Gazette*.

82

² www.canada.ca/en/environment-climate-change/services/species-risk-act-accord-funding.html#2

³ These federally protected areas are: a national park of Canada named and described in Schedule 1 to the *Canada National Parks Act*; The Rouge National Park established by the *Rouge National Urban Park Act*; a marine protected area under the *Oceans Act*; a migratory bird sanctuary under the *Migratory Birds Convention Act, 1994*; or a national wildlife area under the *Canada Wildlife Act*. See ss. 58(2) of SARA.

83 For critical habitat located on other federal lands, the competent minister must either
84 make a statement on existing legal protection or make an order so that the prohibition
85 against destruction of critical habitat applies.

86
87 If the critical habitat for a migratory bird is not within a federal protected area and is not
88 on federal land, within the exclusive economic zone or on the continental shelf of
89 Canada, the prohibition against destruction can only apply to those portions of the
90 critical habitat that are habitat to which the *Migratory Birds Convention Act, 1994* applies
91 as per SARA ss. 58(5.1) and ss. 58(5.2).

92
93 For any part of critical habitat located on non-federal lands, if the competent minister
94 forms the opinion that any portion of critical habitat is not protected by provisions in or
95 measures under SARA or other Acts of Parliament, or the laws of the province or
96 territory, SARA requires that the Minister recommend that the Governor in Council make
97 an order to prohibit destruction of critical habitat. The discretion to protect critical habitat
98 on non-federal lands that is not otherwise protected rests with the Governor in Council.
99

100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120

Acknowledgments

This recovery strategy was prepared by Candace Neufeld (Environment and Climate Change Canada – Canadian Wildlife Service, Prairie Region) with contributions from Sarah Lee (Environment and Climate Change Canada – Canadian Wildlife Service, Prairie Region). Valuable reviews were provided by Yeen Ten Hwang and Medea Curteanu (Environment and Climate Change Canada – Canadian Wildlife Service, Prairie Region), Gina Schalk (Environment and Climate Change Canada – National Capital Region), and Manitoba Conservation.

The Manitoba Conservation Data Centre provided updated element occurrence information; Chris Friesen and Colin Murray provided valuable insight on the species occurrences and habitat requirements in Manitoba. Acknowledgement and thanks is given to all other parties that provided advice and input used to help inform the development of this recovery strategy including various Indigenous organizations and individuals, landowners, citizens and stakeholders who provided input and/or participated in consultations. The co-operation of all the landowners, lessees and land managers who granted access to their land to do surveys and who continue to provide habitat for species at risk is greatly appreciated.

121 **Executive Summary**

122
123 The Fascicled Ironweed (*Vernonia fasciculata*) is a perennial wildflower, producing
124 30-120 cm tall stems, with flat-topped clusters of purple flowers, and lance-shaped
125 leaves alternating up the stem. Fascicled Ironweed flowers in July and August. Seeds
126 are wind-dispersed but are likely also dispersed by flowing water. In Manitoba, plants
127 inhabit moist to wet soils in open to semi-open riparian areas, sedge and wet meadows,
128 river terraces, oxbows, banks of rivers and creeks, low-lying depressions in prairie, and
129 ditches and can tolerate seasonal flooding.

130
131 Fascicled Ironweed is widespread in the tallgrass prairies of midwestern and eastern
132 United States and into southern Manitoba. In Canada, Fascicled Ironweed is designated
133 as Endangered on Schedule 1 of the federal *Species at Risk Act* (SARA). As of 2019,
134 there were two extant populations in Manitoba, along with two historical populations that
135 are likely extirpated (one in Manitoba and one in Saskatchewan). The Canadian
136 population is coarsely estimated at 21,000 plants, although the majority of the plants are
137 in one population (Rat River, MB), with less than 100 plants at the second population
138 (Mile Road 4W, MB).

139
140 Additional loss of habitat quantity or quality among the known populations of Fascicled
141 Ironweed could adversely affect the species' survival in Canada. Threats, in order of
142 highest to lowest threat impact, are: annual and perennial non-timber crops (cultivation);
143 dams and water management/use (alteration of flood duration/frequency); housing and
144 urban areas (residential development and landscaping); agricultural effluents (herbicide
145 use); roads (road construction and maintenance); livestock ranching (soil disturbance by
146 livestock) and invasive non-native plant species.

147
148 Recovery is considered feasible for this species. The population and distribution
149 objectives are to maintain the estimated population size and distribution, within the
150 natural range of variation, of the extant populations as well as to maintain any newly
151 discovered or re-discovered populations, and to ensure long-term persistence and
152 natural expansion of Fascicled Ironweed in Canada. Broad strategies to be taken to
153 address the threats to the survival and recovery of Fascicled Ironweed are presented in
154 the section on Strategic Direction for Recovery.

155
156 Critical habitat is fully identified in this recovery strategy for all extant populations in
157 Canada. In Manitoba, critical habitat is identified as all occupied suitable habitat and all
158 natural biophysical attributes within a 300 m critical function zone extending from the
159 outer boundary of the occupied suitable habitat.

160
161 One or more action plans for Fascicled Ironweed will be posted on the Species at Risk
162 Public Registry within five years following the final posting of this recovery strategy.

163 **Recovery Feasibility Summary**

164
165 Based on the following four criteria that Environment and Climate Change Canada uses
166 to establish recovery feasibility, recovery of the Fascicled Ironweed has been deemed
167 technically and biologically feasible.

168
169 **1. Individuals of the wildlife species that are capable of reproduction are available**
170 **now or in the foreseeable future to sustain the population or improve its**
171 **abundance.**

172
173 Yes. Several thousand Fascicled Ironweed individuals capable of reproduction are
174 present, primarily in the Rat River population in Manitoba. Fascicled Ironweed may
175 slowly expand through vegetative reproduction with underground rhizomes. Flower and
176 seed production have also been observed at the extant populations. Assuming that no
177 other major threats impact the species or its habitat, the Rat River population is
178 expected to sustain itself and persist as it has historically; it is unknown whether the
179 Mile Road 4W population is self-sustaining over the long-term due to recent loss of
180 individuals and habitat through cultivation and ditch maintenance.

181
182 **2. Sufficient suitable habitat is available to support the species or could be made**
183 **available through habitat management or restoration.**

184
185 Yes. Suitable habitat currently exists where extant populations occur and the habitat
186 should be sufficient to maintain or increase species persistence at current levels, with
187 natural population fluctuations. Beneficial management practices have the potential to
188 maintain and enhance the habitat, possibly creating additional suitable habitat within the
189 current distribution. Unoccupied suitable habitat is available in small quantities.
190 Roadside ditches would likely be suitable, given that historical and current populations
191 reside in these habitats, although occurrences inhabiting these areas would be at risk
192 from threats mentioned in section 4.2 and would be quite fragmented. Remnant moist to
193 wet meadows still exist near the known extent of Fascicled Ironweed in Manitoba,
194 primarily in the Manitoba Tall Grass Prairie Preserve, however, the majority of wet
195 prairie in southern Manitoba has been converted to agricultural fields. Additional riparian
196 habitat would likely be suitable (e.g. further downstream or upstream of the main Rat
197 River population or in adjacent tributaries), although many waterways in the area have
198 been channelized or otherwise altered for faster drainage, or riparian areas have been
199 cleared for agriculture or development (COSEWIC 2014, Manitoba Conservation and
200 Water Stewardship unpubl. data).

201
202 **3. The primary threats to the species or its habitat (including threats outside**
203 **Canada) can be avoided or mitigated.**

204
205 Yes. Identified threats are anthropogenic, related to loss in habitat quality and quantity,
206 and can be mitigated through beneficial management practices, habitat stewardship/
207 conservation, increased awareness through communication of the species' needs and

208 threats and incorporating those considerations into land use planning. It should be
209 noted, however, that Fascicled Ironweed occurs at the northern limit of its range in
210 Canada, and is primarily concentrated in one population, making it vulnerable to
211 stochastic events and possibly genetic inbreeding.

212

213 **4. Recovery techniques exist to achieve the population and distribution**
214 **objectives or can be expected to be developed within a reasonable timeframe.**

215

216 Yes. Recovery techniques such as increasing landowner and land manager awareness
217 of the species' habitat requirements through communication and engagement strategies
218 and land use planning, developing and employing adaptive beneficial management
219 practices, using stewardship agreements or conservation easements for habitat
220 conservation, continuing with inventory and monitoring work to assess population
221 health, and conducting research into knowledge gaps can be developed within a
222 reasonable timeframe, given adequate resources, and should contribute to achieving
223 the population and distribution objective.

224

225

226

227	Table of Contents	
228		
229	Preface.....	i
230	Acknowledgments.....	iii
231	Executive Summary	iv
232	Recovery Feasibility Summary.....	v
233	1. COSEWIC Species Assessment Information.....	1
234	2. Species Status Information	1
235	3. Species Information	2
236	3.1 Species Description	2
237	3.2 Species Population and Distribution.....	2
238	3.3 Needs of the Fascicled Ironweed	6
239	4. Threats.....	8
240	4.1 Threat Assessment	8
241	4.2 Description of Threats	10
242	5. Population and Distribution Objectives.....	13
243	6. Broad Strategies and General Approaches to Meet Objectives	15
244	6.1 Actions Already Completed or Currently Underway	15
245	6.2 Strategic Direction for Recovery.....	16
246	6.3 Narrative to Support the Recovery Planning Table	18
247	7. Critical Habitat.....	19
248	7.1 Identification of the Species' Critical Habitat	19
249	7.2 Activities Likely to Result in the Destruction of Critical Habitat.....	21
250	8. Measuring Progress	24
251	9. Statement on Action Plans	24
252	10. References.....	25
253	Appendix A: Summary of Fascicled Ironweed Populations in Canada	31
254	Appendix B: Critical Habitat for Fascicled Ironweed in Canada	33
255	Appendix C: Effects on the Environment and Other Species	35
256		

257
258

1. COSEWIC* Species Assessment Information

Date of Assessment: November 2014

Common Name (population): Fascicled Ironweed

Scientific Name: *Vernonia fasciculata*

COSEWIC Status: Endangered

Reason for Designation: This showy perennial plant has a restricted geographic range in Canada, and occupies small prairie remnants mainly along roadside ditches and riversides in southern Manitoba. The few small subpopulations are at risk from such threats as flood duration/frequency alteration, cultivation, ranching, herbicide use, and road and right-of-way maintenance activities.

Canadian Occurrence: Manitoba

COSEWIC Status History: Designated Endangered in November 2014

259
260
261

* COSEWIC (Committee on the Status of Endangered Wildlife in Canada)

262
263
264
265
266
267
268
269
270

2. Species Status Information

Fascicled Ironweed (*Vernonia fasciculata*) is designated as Endangered on Schedule 1 of the federal *Species at Risk Act* (SARA), and as Endangered under Manitoba's *Endangered Species and Ecosystems Act*. The conservation status of Fascicled Ironweed throughout its range is described in Table 1. It is estimated that the Canadian range is less than 1% of the species' global range.

Table 1. NatureServe conservation status of Fascicled Ironweed (NatureServe 2020b)^a.

Global (G) Rank ^b	National (N) Rank ^b	Sub-national (S) Rank ^b
G5	Canada (N1)	Manitoba (S1), Saskatchewan (SH)
	United States (N5?) ^c	Colorado (SNR), Illinois (SNR), Indiana (SNR), Iowa (S5), Kansas (SNR), Massachusetts (SNR), Michigan (SNR), Minnesota (SNR), Mississippi (SNR), Missouri (SNR), Nebraska (SNR), New York (SNR), North Dakota (SNR), Ohio (S2), Oklahoma (SNR), South Dakota (SNR), Wisconsin (SNR)

271
272
273
274

^a Two subspecies of *Vernonia fasciculata* have been described (subspecies *corymbosa* and subspecies *fasciculata*) and are included in NatureServe (2020b) and in Flora of the Great Plains (Great Plains Flora Association 1986). However, COSEWIC (2014), the Biotic of North American Program (BONAP; Kartesz 2015), and the Flora of North America (Strother 2006) only recognize *Vernonia fasciculata* to the species

275 level, with both describing *Vernonia fasciculata* ssp *corymbosa* as a synonym of *Vernonia fasciculata*.
276 Therefore, the conservation status listed here will be to the species level only, and will include distribution
277 notes from other sources below in the footnotes.
278 ^b The NatureServe conservation status of a species is designated by a number from 1 to 5, preceded by a
279 letter reflecting the appropriate geographic scale of the assessment (G = Global, N = National, and
280 S = Subnational). The numbers have the following meaning: 1 = critically imperiled, 2 = imperiled,
281 3 = vulnerable, 4 = apparently secure, and 5 = secure. NR = not ranked, ? = inexact or uncertain and
282 qualifies the character immediately before it (NatureServe 2020c).
283 ^c COSEWIC (2014) includes Kentucky (Jones 1972, Medley 1993, Strother 2006), Arkansas (Smith 1973,
284 1988), Alabama (no source), Montana (Lesica et al. 1984, Great Plains Flora Association 1986) and
285 Texas (White 2012) in the global distribution map, as does Kartesz (2015) with the exception of Alabama
286 and Arkansas, and USDA (2019) with the exception of Alabama, but NatureServe (2020b) does not
287 include these states, nor does the Flora of North America (Strother 2006) with the exception of Kentucky.
288 The Montana government states Fascicled Ironweed does not occur in Montana due to there being no
289 herbarium record or relocation of this species, and believe the inclusion of this species in Montana in the
290 Flora of the Great Plain (Great Plains Flora Association 1977) to be in error (Montana Natural Heritage
291 Program 2019). NatureServe (2020b) includes a conservation status for New York and Massachusetts
292 but COSEWIC (2014) and Kartesz (2015) list those as introduced populations. Kartesz (2015) and Flora
293 of North America (Strother 2006) do not include Mississippi in the global distribution map but a herbarium
294 specimen from Mississippi is located in the Delta State University Herbarium.
295
296

297 **3. Species Information**

298 299 **3.1 Species Description**

300
301 Fascicled Ironweed is a perennial wildflower which grows upright to 30-120 cm tall
302 either as single stems or several stems clustered around a tough, fibrous-rooted base
303 (Great Plains Flora Association 1986). Its stem is sometimes reddish at the base with
304 many lance-shaped, stalkless leaves, alternating evenly up the stem; leaves have
305 pointed teeth around the edge and small pits on the underside containing awl-shaped
306 hairs (Great Plains Flora Association 1986). Plants flower in July and August. The
307 inflorescence is a wide, flat-topped cluster of flower heads, each made up of 10-26
308 purple tubular (disc) florets (see cover photo). Plants spread by seeds which disperse
309 by wind, due to the bristles (pappus) attached to the seeds (achenes), but likely also
310 disperse by flowing water, as well as vegetatively by means of horizontal underground
311 stems (rhizomes).
312

313 **3.2 Species Population and Distribution**

314 315 ***Global Distribution***

316
317 Fascicled Ironweed is native to North America, occurring in both Canada and the United
318 States. In Canada, its northern extent is limited to tallgrass prairie remnants in southern
319 Manitoba, although a historical record exists from southeastern Saskatchewan. In the
320 United States, it is present in central and mid-eastern States in the tallgrass prairie
321 (Table 1, Fig. 1). Population data is not available for Fascicled Ironweed across its
322 North American range.
323



324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339

Figure 1. Current distribution of Fascicled Ironweed in North America (adapted from Strother 2006, COSEWIC 2014, Kartesz 2015, NatureServe 2020b, USDA 2019). The black circle indicates the historical and likely extirpated record from Saskatchewan and the question marks indicate questionable records (see Table 1).

340 **Canadian Distribution**

341
 342 As of March 2019 in Canada, there were two⁴ known extant⁵ native populations⁶ in
 343 Manitoba (Fig. 2, Appendix A). Both populations are in the Prairie Ecozone, and the
 344 Lake Manitoba Plain Ecoregion. The largest population (Rat River) occurs adjacent to,
 345 or near, the Rat River. The second population (Mile Road 4W) is now limited to a small
 346 patch in a ditch near Lowe Farm, after the adjacent native prairie where the majority of
 347 the plants had resided was cultivated and the ditch was deepened. The Canadian
 348 population is estimated at 21,000⁷ plants, with an extent of occurrence⁸ of 338 km² and
 349 an index area of occupancy of 60 km² (COSEWIC 2014, Manitoba Conservation Data
 350 Centre unpublished data 2019, Manitoba Conservation Data Centre personal
 351 communication 2019). The actual area of occupancy⁹ of the population has never been
 352 mapped so the full extent of the population along the Rat River, particularly as it extends
 353 away from the shore line, is not known. There has also never been an inventory done
 354 on the same year of the entire Rat River population to obtain a more accurate and
 355 precise estimate of population size; the actual counts of plants based on portions of the
 356 population are considerably less than the estimated overall population size (Table A1 in
 357 Appendix A).

358
 359 Details on the original distribution and abundance of Fascicled Ironweed in Canada are
 360 not known (COSEWIC 2014). Although Fascicled Ironweed may always have been rare
 361 at its northern extent, it is probable it would have been more prevalent in Manitoba and
 362 possibly Saskatchewan, prior to habitat conversion from agricultural activities and
 363 settlements. Historical records exist from the Otterburne and Morris areas of Manitoba;
 364 the historical record from Morris area is likely extirpated (Table A1 in Appendix A, and

⁴ COSEWIC (2014) lists “Provincial Road (PR) 200” and “Rat River” as separate element occurrences, but additional surveys found Fascicled Ironweed further down the Rat River closer to PR 200, which merged the two into one element occurrence based on NatureServe (2020a) habitat-based element occurrence delimitation guidance.

⁵ Extant means the population has been recently verified as still existing, information is accurate, and habitat still exists at the time of writing.

⁶ For the purposes of this recovery strategy, a population will be considered equivalent to an element occurrence as defined by NatureServe (2020a) and the Manitoba Conservation Data Centre and a subpopulation as defined by COSEWIC (2014). Populations may be comprised of one or more occurrences (patches of plants).

⁷ From the COSEWIC (2014) status report regarding the population estimate of the Rat River population: “Considering the number of patches consisting of 1,000 or more stems (sometimes many more than 1,000), plus the number of smaller patches, it does not seem unreasonable to coarsely estimate the number of stems along the Rat River to be 125,000 ± 25,000 (20,833 plants ± 4,167 plants). Given the lack of precise estimation of the number of plants, a relatively high uncertainty value (± 25,000 stems) was chosen. This assessment will use the rounded value of 21,000 as the estimated number of plants in the Rat River subpopulation, but this is a very coarse estimate.” COSEWIC (2014) used the median number of stems per plant (six) to estimate number of plants.


⁸ Extent of occurrence and index area of occupancy are as defined by COSEWIC 2020a and COSEWIC 2020b, respectively.

⁹ Area of occupancy, as defined in the recovery strategy, is the actual area on the ground occupied by Fascicled Ironweed plants. Area of occupancy is often determined by walking the perimeter of the patches of plants using a GPS unit.

365 see Table 1 in COSEWIC 2014). There is one historical record from Saskatchewan in
 366 the Weyburn area, observed sometime prior to 1966, but that has never been relocated
 367 despite targeted search effort and is likely extirpated (Table A1 in Appendix A).

368
 369 Determining trends in population size of Fascicled Ironweed in Canada are not possible
 370 for a few reasons. This species has not been consistently monitored, with portions of
 371 the primary population along the Rat River being visited only once. Targeted surveys
 372 with estimates of plant counts only began in 2005. Fluctuations of numbers of plants
 373 appears to happen with moisture availability although it isn't clear whether the number
 374 of plants is actually changing or whether it is an issue with detection of plants between
 375 dry and wet years (COSEWIC 2014).

376



377
 378 **Figure 2.** Current distribution of Fascicled Ironweed in Canada (compiled from Manitoba
 379 Conservation Data Centre data 2019).

380

381

382

383

384

385

386

387 3.3 Needs of the Fascicled Ironweed

388
389 In Manitoba, Fascicled Ironweed is now confined to remnant native habitat that has not
390 been cultivated, or ditches and right of ways. It is unknown what the habitat and
391 distribution would have been prior to European settlement and cultivation but it is likely
392 quite restricted now compared to its original range given the amount of decline in moist
393 prairie meadows and riparian forest habitat (COSEWIC 2014, Murray and Church
394 2017). In other parts of its North American range, it typically occupies wet meadows,
395 banks of creeks and wetlands, roadsides/ditches, floodplains, and low-lying moist
396 depressions or swales in prairie, often reported on sandy soils, but is found occasionally
397 in the drier upland sandhill prairie (Strother 2006, Consortium of Pacific Northwest
398 Herbaria unpubl. data 2015, Kansas State University Herbarium unpubl. 2017).

399
400 In Manitoba to date, the largest population grows in moist to wet clay soils in open to
401 semi-open (deciduous treed) riparian areas including moist to wet meadows, river
402 terraces, oxbows, banks of rivers and creeks and low-lying depressions, primarily along
403 the margins of the Rat River (Foster and Reimer 2007, COSEWIC 2014, Manitoba
404 Conservation unpubl. data 2019). The habitat along the Rat River is subject to seasonal
405 flooding (Foster and Reimer 2007, COSEWIC 2014). Fascicled Ironweed has been
406 recorded within 40-100 m of the Rat River, although surveys have often not extended
407 into the upland beyond that distance; plants are in a band between the riverbank and
408 the riparian forest or cultivated upland (Murray and Friesen 2012, Manitoba
409 Conservation Data Centre unpubl. data 2019). The semi-open treed areas have an
410 overstory of native tree species while open and semi-open areas have a very sparse to
411 dense herbaceous understory containing both native and non-native plant species¹⁰
412 (Foster and Reimer 2007, Manitoba Conservation Data Centre unpubl. data 2019).
413 Fascicled Ironweed is sometimes the dominant or co-dominant species in the
414 understory along the Rat River (Foster and Reimer 2007). Fascicled Ironweed has not
415 been found growing in areas along the bank of the Rat River where there was very
416 thick, dense vegetation from grasses and sedges with little to no tree cover, nor does it
417 appear to tolerate dense shade (COSEWIC 2014, Manitoba Conservation Data Centre,
418 unpubl. data 2019). In a few places along the Rat River, and at the Mile Road 4W
419 population, Fascicled Ironweed grows in ditches; the ditches contain some introduced
420 species of grass (COSEWIC 2014, Manitoba Conservation Data Centre unpubl. data
421 2019).

422

¹⁰Tree species recorded in the overstory include Green Ash (*Fraxinus pennsylvanica*), Manitoba Maple (*Acer negundo*), American Elm (*Ulmus americanus*), Bur Oak (*Quercus macrocarpa*), Eastern Cottonwood (*Populus deltoides*) and willow species (*Salix* species). Native herbaceous species recorded in the understory include Tall Beggarticks (*Bidens vulgata*), Hedge Bindweed (*Calystegia sepium*), Riverbank Grape (*Vitis riparia*), Wild Cucumber (*Echinocystis lobata*), Carrion flower (*Smilax lasioneura*), Canada Moonseed (*Menispermum canadense*), Poison Ivy (*Toxicodendron rydbergii*), Wild Mint (*Mentha arvensis*), Alkali Cordgrass (*Spartina gracilis*), Slender Wildrye (*Elymus trachycaulus*) and Smooth-fruited Sedge (*Carex laeviconica*) among others, while non-native species include Narrow-leaved Dock (*Rumex stenophyllus*), Smooth Brome (*Bromus inermis*), Reed Canarygrass (*Phalaris arundinaceaea*), Sow Thistle (*Sonchus arvensis*), Common Plantain (*Plantago major*), and Common Hemp-nettle (*Galeopsis tetrahit*) (Foster and Reimer 2007, Manitoba Conservation Data Centre unpubl. data 2019).

423 *Limiting Factors*

424

425 Fascicled Ironweed reaches the northern extreme of its range in Canada. Populations at
426 the limit of a species' range often occupy poorer habitat and are more fragmented, less
427 dense, and more variable than those at their core range (Channell and Lomolino 2000,
428 Vucetich and Waite 2003). These peripheral populations are therefore more vulnerable
429 to extinction due to low immigration rate, disrupted pollinator relationships, and other
430 density-related factors. Genetic diversity is sometimes, but not always, less in peripheral
431 populations, but they may possess unique genetic characteristics (Vucetich and Waite
432 2003). Since there is only one population in Manitoba containing almost all of the
433 Fascicled Ironweed plants, and this population is greatly isolated from populations in the
434 United States, it may make it more vulnerable to stochastic events (e.g. extreme or
435 prolonged flooding events) and genetic issues like inbreeding depression, due to
436 isolation from other populations , although research into this is needed.

437

438 **4. Threats**

439
440 **4.1 Threat Assessment**

441
442 The Fascicled Ironweed threat assessment is based on the IUCN-CMP (World Conservation Union–Conservation
443 Measures Partnership) unified threats classification system. Threats are defined as the proximate activities or processes
444 that have caused, are causing, or may cause in the future the destruction, degradation, and/or impairment of the entity
445 being assessed (population, species, community, or ecosystem) in the area of interest (global, national, or subnational).
446 Limiting factors are not considered during this assessment process. For purposes of threat assessment, only present and
447 future threats are considered. Historical threats, indirect or cumulative effects of the threats, or any other relevant
448 information that would help understand the nature of the threats are presented in the Description of Threats section.

449 **Table 2.** Threat calculator assessment.
450

Threat #	Threat description	Impact ^a	Scope ^b	Severity ^c	Timing ^d	Detailed threats
1	Residential & commercial development	Low	Small	Extreme-Moderate	Moderate	
1.1	Housing & urban areas	Low	Small	Extreme-Moderate	Moderate	Acreages, farm yards, rural lots with landscaping or mowing up to the river's edge
2	Agriculture & aquaculture	Medium-Low	Restricted-Small	Extreme-Serious	Moderate	
2.1	Annual & perennial non-timber crops	Medium-Low	Restricted-Small	Extreme-Serious	Moderate	Cultivation
2.3	Livestock farming & ranching	Negligible	Negligible	Serious-Slight	Moderate	Soil disturbance by livestock
4	Transportation & service corridors	Negligible	Negligible	Extreme-Serious	High	
4.1	Roads & railroads	Negligible	Negligible	Extreme-Serious	High	Road and ditch maintenance and construction

Threat #	Threat description	Impact ^a	Scope ^b	Severity ^c	Timing ^d	Detailed threats
7	Natural system modifications	Medium-Low	Pervasive	Moderate-Slight	High	
7.2	Dams & water management/use	Medium-Low	Pervasive	Moderate-Slight	High	Alteration of flood duration/frequency from dams, drains, dykes, diversions and other flood control measures
8	Invasive & other problematic species & genes	Negligible	Negligible	Unknown	High	
8.1	Invasive non-native/alien species	Negligible	Negligible	Unknown	High	Invasive non-native species
9	Pollution	Low	Restricted	Moderate	High	
9.3	Agricultural & forestry effluents	Low	Restricted	Moderate	High	Herbicide use in adjacent fields

451 ^a **Impact** – The degree to which a species is observed, inferred, or suspected to be directly or indirectly threatened in the area of interest. The
452 impact of each threat is based on Severity and Scope rating and considers only present and future threats. Threat impact reflects a reduction of a
453 species population or decline/degradation of the area of an ecosystem. The median rate of population reduction or area decline for each
454 combination of scope and severity corresponds to the following classes of threat impact: Very High (75% declines), High (40%), Medium (15%),
455 and Low (3%). Unknown: used when impact cannot be determined (e.g., if values for either scope or severity are unknown); Not Calculated:
456 impact not calculated as threat is outside the assessment timeframe (e.g., timing is insignificant/negligible or low as threat is only considered to be
457 in the past); Negligible: when scope or severity is negligible; Not a Threat: when severity is scored as neutral or potential benefit.

458 ^b **Scope** – Proportion of the species that can reasonably be expected to be affected by the threat within 10 years. Usually measured as a
459 proportion of the species' population in the area of interest. (Pervasive = 71–100%; Large = 31–70%; Restricted = 11–30%; Small = 1–10%;
460 Negligible < 1%).

461 ^c **Severity** – Within the scope, the level of damage to the species from the threat that can reasonably be expected to be affected by the threat
462 within a 10-year or three-generation timeframe. Usually measured as the degree of reduction of the species' population. (Extreme = 71–100%;
463 Serious = 31–70%; Moderate = 11–30%; Slight = 1–10%; Negligible < 1%; Neutral or Potential Benefit ≥ 0%).

464 ^d **Timing** – High = continuing; Moderate = only in the future (could happen in the short term [< 10 years or 3 generations]) or now suspended
465 (could come back in the short term); Low = only in the future (could happen in the long term) or now suspended (could come back in the long
466 term); Insignificant/Negligible = only in the past and unlikely to return, or no direct effect but limiting.

4.2 Description of Threats

Appendix A identifies the threats associated with each population. Threats are discussed below in decreasing order of Level 1 threat impact.

IUCN Threat 2. Agriculture and Aquaculture

Threat 2.1 Annual & perennial non-timber crops

The threat of cultivation is mostly historical. Historical conversion of upland tallgrass prairie and adjacent riparian habitats to cultivated cropland likely contributed to the loss of Fascicled Ironweed habitat (Murray and Friesen 2012, COSEWIC 2014). Almost all suitable upland habitat has been cultivated along the Rat River population (EO 2709), sometimes right up to the river's edge. The population at Mile Road 4W (EO 4867) declined significantly when the prairie upland was converted to agricultural land, leaving only a small portion of the population remaining in the ditch (Foster and Reimer 2007, Friesen and Murray 2011, COSEWIC 2014). There are portions of uncultivated upland habitat remaining along the Rat River, some of which would not be suitable for agriculture due to flooding, poor grade, or moist conditions but which may be cultivated in drought conditions. There are also areas along the Rat River where Fascicled Ironweed is growing immediately adjacent to agricultural fields, and are at risk of cultivation. However, as the majority of the Rat River population occurs within 40 m from the water, the impact of encroaching cultivation on the Fascicled Ironweed population as a whole is limited.

Threat 2.3 Livestock farming & ranching

Fascicled Ironweed is suspected to be unpalatable to cattle despite having good nutritional value and crude protein levels (Hubbard and Boe 1988), and has been reported to increase in wet pastures where cattle graze around it (Shaw and Schmidt 2003). COSEWIC (2014) reported Fascicled Ironweed absent or declining in an area along the Rat River heavily grazed by cattle but present adjacent to the grazed area; the absence of plants in the grazed area may be a result of trampling or other soil disturbance caused by the cattle.

IUCN Threat 7. Natural System Modifications

Threat 7.2 Dams & water management/use

Fascicled Ironweed appears to be flood tolerant (Shaw and Schmidt 2003). Its seeds are buoyant due to bristly hairs (pappi) attached to the seed, and they likely disperse downstream through the water (hydrochory) or further into the upland during times of high flow or floods (Groves 2010, Catford et al. 2014, Carthey et al. 2016). In areas where it floods regularly, such as along the Rat River, Fascicled Ironweed likely has a competitive advantage over plants that aren't flood tolerant due to its ability to spread through rhizomes (Sluis and Tandarich 2004, Catford et al. 2014); these rhizomes may

513 also break off and drift downstream and form new plants if they root in suitable habitat.
514 In addition, seasonal floods or higher water levels likely maintain the wet meadows and
515 riparian areas that are required habitat for Fascicled Ironweed. Upstream of the Rat
516 River population (EO 2709), there is a major dam at St. Malo along with smaller dams,
517 drains, dykes, diversions etc. These water control structures are in an attempt to control
518 floods and water levels along the Rat River. Lowering of water levels or changes in the
519 duration and frequency of floods through dams, diversions and flow regulation
520 structures may result in a decline in habitat suitability for seed germination and seedling
521 establishment for Fascicled Ironweed or change the plant community composition along
522 the Rat River (Jansson et al. 2005, Uowolo et al. 2005, Merritt and Wohl 2006). Flow
523 regulation structures in rivers can also physically prevent dispersal of seeds
524 downstream, or affect how far seeds disperse and where they are deposited (Merritt
525 and Wohl 2002, Brown and Chenoweth 2008). Since more than 99% of the Fascicled
526 Ironweed in Manitoba resides along the Rat River, this threat is pervasive in scope.

527

528 **IUCN Threat 1. Residential and Commercial Development**

529

530 *Threat 1.1 Housing & urban areas*

531

532 There are over 50 farm yards or acreages, in addition to the community of Otterburne,
533 along the stretch of Rat River containing Fascicled Ironweed (EO 2709). Some of these
534 parcels of land have mowed lawn and landscaping where there would naturally have
535 been wet meadows or riparian treed areas; this existing activity is considered historical
536 and not included in the scope of this threat assessment. The remaining parcels are
537 either cultivated very close to the river's edge (threat 2.1) or have a buffer of varying
538 width of riparian vegetation, trees, and/or wet meadow. These areas are mostly
539 unsuitable for housing development due to annual flood risk and poor grade but
540 landowners may alter the remaining natural areas by landscaping (e.g. removing natural
541 vegetation, grading the area, putting in non-native lawn grass) or regular mowing as
542 others have done, particularly with successive drought years. Depending on the extent
543 of the activity, it could alter the habitat and destroy Fascicled Ironweed plants growing in
544 these areas. There is also a golf course bordering the Rat River, but since it has already
545 been created, it is considered a historical threat; if there were plans to expand the golf
546 course, this threat would need to be re-evaluated.

547

548 **IUCN Threat 9. Pollution**

549

550 *Threat 9.3 Agricultural & forestry effluents*

551

552 Use of pesticides intended to control undesirable plants (broad-leaf herbicide) or insect
553 pests (indirectly through herbicide or directly through insecticide) in areas occupied by,
554 or adjacent to, Fascicled Ironweed can affect Fascicled Ironweed plants and its
555 pollinators. Broad-leaf herbicides directly sprayed on Fascicled Ironweed will kill the
556 plant, and use of broad-leaved herbicides in fields adjacent to Fascicled Ironweed may
557 drift during application or leach out with rains and damage or kill Fascicled Ironweed
558 plants. Targeted spot applications or other means of invasive non-native species control

559 are required in areas with Fascicled Ironweed plants. Fascicled Ironweed occurs directly
560 adjacent to agricultural fields, yards, and golf courses, as well as in ditches in some
561 areas along the Rat River and at Mile Road 4W, and is therefore at risk from improper
562 use of pesticides in those areas (Murray and Church 2015).

563
564 In general, reducing flowering plants in an area through herbicide use, as well as
565 spraying insecticides to control insect pest species, can both reduce pollinators and
566 potentially affect seed production of those plants. Fascicled Ironweed relies primarily on
567 insect pollinators such as long-tongued bees (Apidae, Anthophoridae, Megachilidae),
568 and short-tongued bees (Halictinae), but also bee flies (Bombyliidae), ants (Formicidae),
569 butterflies (Nymphalidae, Lycaenidae, Pieridae, Papilionidae), skippers (Hesperiidae),
570 moths (Noctuidae) and beetles (Cantharidae) for pollination (Reed 1993, Discover Life
571 2019, Hilty 2019). Declining native bee populations across North America have been
572 observed to coincide with declines in native plant populations, although it is not clear
573 whether bee declines are causing plant declines or vice versa (Nabhan and Buchmann
574 1997, Kearnes et al. 1998, Gill and Raine 2014, Godfray et al. 2014, Scheper et al.
575 2014). However, there is emerging evidence in the literature that pesticide use in
576 croplands is one of the major factors causing bee declines worldwide either through
577 acute (lethal doses causing death) or chronic effects (sublethal doses altering behavior)
578 (Gill and Raine 2014, Godfray et al. 2014).

579
580 As of 2014, neonicotinoids were being used on more than 40 million hectares of
581 cropland in the United States and are now the most widely used insecticide in the world,
582 trends that are consistent with the Canadian use of this insecticide (Douglas and Tooker
583 2015). Research has shown that even sublethal exposure to this insecticide can cause
584 acute and chronic effects in pollinating species, specifically in social bees (honeybees,
585 bumblebees, and stingless bees) (Gill and Raine 2014, Godfray et al. 2014). Chronic
586 exposure to sublethal doses of neonicotinoids can alter bumblebee (*Bombus spp.*)
587 behavior including changing forager preferences for flower types, impairing forager
588 performance (carry out fewer foraging bouts and bring back smaller pollen loads), and
589 impair bee learning performance (Gill and Raine 2014, Godfray et al. 2014). Bees are
590 the main pollinator of Fascicled Ironweed and effects from changes in bee behaviour
591 could be exacerbated by the competition for pollinators with other co-flowering plants.
592 Further research into this and its potential impact on Fascicled Ironweed is needed.

593

594 **IUCN Threat 4. Transportation and Service Corridors**

595

596 *Threat 4.1 Roads & railroads*

597

598 Habitat, plants, and seed banks can be damaged or destroyed by road construction or
599 maintenance activities such as road widening, realigning or improving the road, ditch
600 deepening/widening (cleaning), trenching, and drainage projects. Habitat and plants can
601 also be affected by incompatible or inappropriately-timed road maintenance activities on
602 shoulders and in ditches, such as spraying pesticides (threat 9.3), grading, haying or
603 mowing. Portions of the Fascicled Ironweed population along the Rat River (EO 2709)
604 are growing where roads cross the river. A portion of this population is adjacent to the

605 Rat River in a roadside ditch and drain which has been mowed in the past with no
606 Fascicled Ironweed plants observed that year (COSEWIC 2014, Manitoba Conservation
607 Data Centre unpubl. data 2019). The remnant population at Mile Road 4W (EO 4867) is
608 in a ditch and at risk from road maintenance activities (Murray and Friesen 2012). It was
609 almost eliminated after the ditch was cleared of vegetation for faster drainage
610 (COSEWIC 2014, Manitoba Conservation Data Centre unpubl. data 2019).

611

612 **IUCN Threat 8. Invasive and Other Problematic Species and Genes**

613

614 *Threat 8.1 Invasive non-native/alien species*

615

616 In Manitoba, Fascicled Ironweed often co-occurs and competes with "weedy" non-native
617 species due to its preference for disturbed habitats where non-native species have an
618 easier time getting established (Shaw and Schmidt 2003, Foster and Reimer 2007).
619 Fascicled Ironweed can have a weedy growth habit and be quite competitive with other
620 native species or non-native species (Shaw and Schmidt 2003, Foster and Reimer
621 2007). Fascicled Ironweed hasn't always occupied habitat with non-native species,
622 however, and the long-term impacts of this increased competition on its growth,
623 reproductive output, recruitment and survival has not been studied. Despite the
624 competitive ability of Fascicled Ironweed, it was found to be outcompeted and displaced
625 by Reed Canary Grass over a 15 year period in a riparian area in Wisconsin when water
626 levels were lower than normal (Barnes 1999). In particular, there is an introduced non-
627 native strain of Reed Canary Grass which is more aggressive than the native strain and
628 can be an issue due to its large size, dense colonial growth, and ability to quickly
629 colonize newly disturbed areas created by changing water levels (Barnes 1999
630 Lavergne and Molofsky 2007). Reed Canary Grass (strain unknown) was reported
631 growing with Fascicled Ironweed at a location along the Rat River where the population
632 of Fascicled Ironweed was estimated at 10,000 stems or more (Foster and Reimer
633 2007); this area has not been monitored to date (Colin Murray, pers. comm. 2019) so
634 the impact of the Reed Canary Grass on the Fascicled Ironweed is not known at this
635 time, but if aggressive, a portion of the Fascicled Ironweed population could be
636 displaced. Narrow-leaved Dock was also mentioned as a potential concern by Foster
637 and Reimer (2007) as it is can also be quite competitive although its ability to compete
638 with Fascicled Ironweed is unknown.

639

640

641 **5. Population and Distribution Objectives**

642

643 The population and distribution objectives for Fascicled Ironweed are to maintain the
644 estimated population size and distribution, within the natural range of variation, of the
645 extant populations as well as any newly discovered¹¹ populations, to ensure long-term
646 persistence and natural expansion of Fascicled Ironweed in Canada.

647

¹¹ Note that occurrences or populations that are considered historical are excluded from these objectives until such time as they are reconfirmed.

648 In Canada, Fascicled Ironweed is at the northern limit of its range restricted to two
649 populations; there is one large population (Rat River) which contains almost all of the
650 individuals in Canada, and a small population (Mile Road 4W) which has declined due
651 to recent habitat loss and may not be viable (Table A1 in Appendix A). Much of the
652 information known about its presence in Manitoba has come in the last decade with
653 increased survey effort. Substantial increases in the number of populations, index area
654 of occupancy, or extent of occurrence are not likely to be documented in the future
655 given that:

- 656 1) the suitable habitat for Fascicled Ironweed in Manitoba is now limited,
657 fragmented, and declining in quality and quantity,
- 658 2) the new occurrences recently documented have all been within the same
659 population,
- 660 3) a considerable amount of suitable habitat has been surveyed already, and
661 4) the species is at the northern limit of its range and may always have been
662 rare.

663 The known area of occupancy (and index area of occupancy) of the main population
664 along the Rat River should continue to increase incrementally with continued survey
665 effort and mapping of plants in adjacent upland habitat. It is possible additional
666 populations will be found with future survey effort in suitable remaining habitat in
667 Manitoba, particularly upstream or downstream of the Rat River population including
668 river systems connected to the Rat River.

669
670 At this time, it is problematic to define specific quantitative population objectives
671 because of the very coarse estimate of population size, absence of long-term data on
672 population trends, and unknown impacts of year to year fluctuations in population size
673 to indicate the range of natural variability for Fascicled Ironweed populations (section
674 3.2). Similarly, since the full extent of the main Fascicled Ironweed population is not
675 known, setting specific quantitative distribution objectives is also difficult. Employing
676 standardized methodology to obtain an understanding of the full extent (area of
677 occupancy) and density (population size) of the main Fascicled Ironweed population is
678 required, as is collecting multiple years of data on portions of the population to quantify
679 the natural range of variation. Once this information is obtained, more specific targets
680 for population and distribution objectives may be defined.

681
682 For this species to be downlisted to Threatened under the COSEWIC status
683 assessment criteria for category B¹² (small distribution range and decline or fluctuation),
684 additional populations would need be confirmed to increase the number of locations (as
685 defined by COSEWIC 2015), and the observed decline in the extent of occurrence,
686 index area of occupancy, quality of habitat, number of locations, and number of mature
687 individuals would have to cease or be reversed (COSEWIC 2014, 2015). For the
688 reasons bulleted above, the likelihood of the species being downlisted under these
689 criteria is low. Therefore, the population and distribution objective has been set in the

¹² COSEWIC (2014) assessed Fascicled Ironweed as Endangered under the following criteria: small extent of occurrence; small index area of occupancy; known to exist at under 5 locations; and continuing observed decline in extent of occurrence, index area of occupancy, area/extent/quality of habitat, and number of locations or populations – B1ab(i,ii,iii,iv)+2ab(i,ii,iii,iv).

690 context of maintaining the populations and their distribution (index area of occupancy
691 and extent of occurrence), and preventing or reversing further declines in quality and
692 quantity of habitat through beneficial management practices, stewardship agreements
693 and communication strategies with targeted groups (Table 3).
694
695

696 **6. Broad Strategies and General Approaches to Meet** 697 **Objectives**

698

699 **6.1 Actions Already Completed or Currently Underway**

700

701 **Inventory and Monitoring**

702

703

704

705

706

707

708

709

710

711

712

712 **Habitat Assessment, Management and Stewardship**

713

714

- Fascicled Ironweed is listed under Manitoba's *Endangered Species and Ecosystem Act* as Endangered.

715 **6.2 Strategic Direction for Recovery**

716

717 **Table 3.** Recovery Planning Table

Threat or Limitation	Priority ^a	Broad Strategy to Recovery	General Description of Research and Management Approaches
All threats	High	Communication, Collaboration and Engagement	<ul style="list-style-type: none"> • Develop and deliver a communication/outreach strategy targeting landowners and land managers whose properties contain Fascicled Ironweed to raise awareness of the species and its habitat needs and threats (e.g. mowing, cultivation, herbicides, landscaping). • Develop and deliver a communication strategy targeting road maintenance personnel (municipal and provincial) and municipal planners to minimize or eliminate habitat deterioration or destruction during road and ditch maintenance or construction activities. • Develop and deliver a communication strategy targeting provincial, municipal and conservation districts involved with dams and water control in the vicinity of the main Fascicled Ironweed population along the Rat River to discuss impact of hydrological changes on the species. • Encourage land owners and the public to report sightings of Fascicled Ironweed.
All threats	High	Habitat Assessment, Management and Stewardship	<ul style="list-style-type: none"> • Using adaptive habitat management, develop and implement beneficial management practices (BMPs) for the species and its habitat, targeting reduction or mitigation of threats (e.g., dams and water control structures, acreage and residential development, landscaping practices like mowing or seeding non-native species (including grass species that create a dense cover), creating natural buffers at river's edge safe from cultivation, indiscriminate application of herbicides, control of invasive non-native species, using compatible grazing practices, etc.). Monitor effectiveness of BMPS at improving habitat and species presence; amend as necessary. • Mitigate the impact of threats to populations and habitat by engaging landowners and land managers in stewardship agreements, including conservation agreements if feasible,

Threat or Limitation	Priority ^a	Broad Strategy to Recovery	General Description of Research and Management Approaches
			aimed at implementing BMPs and protecting critical habitat. Monitor effectiveness of agreements in conserving habitat. <ul style="list-style-type: none"> • Integrate habitat management with that for other species occurring in the same habitat and surrounding management area (Appendix C). • Promote consistent enforcement or implementation of existing protection measures and regulations (<i>Manitoba Endangered Species and Ecosystems Act</i>).
Knowledge Gaps, all threats	Medium	Inventory and Monitoring	<ul style="list-style-type: none"> • Using standardized protocols (e.g. Henderson 2010a), determine area of occupancy of each Fascicled Ironweed population (element occurrence) and obtain a more precise estimate of population size. • Establish regular monitoring program for Fascicled Ironweed populations, including habitat quality and threats. • Using consistent survey techniques (e.g. Henderson 2010a), continue inventories of suitable habitat to locate new populations; habitat models may be useful in predicting priority search areas (e.g., habitat suitability and/or species distribution models).
Knowledge gaps, all threats	Medium	Research as part of an adaptive management framework	<ul style="list-style-type: none"> • Determine effect of population size and isolation on genetic diversity and population viability, and develop a seed gene bank if deemed necessary. • Conduct research to develop a better understanding of the species ecology and habitat needs (e.g., seed bank viability, seed viability in flooding conditions, impact of hydrochory, recruitment and survival rates, pollination biology, pesticide use on pollinators, genetic exchange) • Assess short- and long-term impacts of threats to Fascicled Ironweed and its habitat quality (e.g. long term impacts of non-native species on Fascicled Ironweed growth and survival, habitat changes with differing water levels). • Apply findings to develop or refine BMPs for the species.

718 ^a “Priority” reflects the degree to which the broad strategy contributes directly to the recovery of the species or is an essential precursor to an
 719 approach that contributes to the recovery of the species

6.3 Narrative to Support the Recovery Planning Table

720
721
722 Due to the continued loss of habitat quality and quantity, and the limited number of
723 populations, engagement with landowners and land managers and encouraging
724 conservation through stewardship are essential to the recovery of Fascicled Ironweed.
725 Habitat requirements of Fascicled Ironweed on private and public lands should be
726 considered during land use planning at all levels (local, municipal, regional, provincial)
727 and during maintenance or construction activities to ensure that land management
728 practices benefitting the species can be implemented. This will be particularly important
729 for occurrences at risk from threats like cultivation, residential development and
730 landscaping, ditch maintenance and road construction, and considerations regarding
731 dams and water control structures. Development and implementation of adaptive
732 site-specific best management practices for the species and its habitat to reduce or
733 mitigate threats from practices such as incompatible landscaping practices for
734 residential lots (e.g. mowing during flowering or seed set, planting non-native species
735 including lawn grass, not leaving a natural buffer at the river's edge), incompatible
736 grazing (trampling along the river's edge), and indiscriminant spraying of pesticides
737 adjacent to cropland is required for successful conservation and may be possible to
738 implement through stewardship agreements.

739
740 Research into aspects of Fascicled Ironweed ecology and impact of its threats is a
741 medium priority but will help with better understanding the habitat and species' needs,
742 and will help refine beneficial management practices. Knowledge gaps exist in areas
743 like seed bank viability, seed viability in flooding conditions, importance of water versus
744 wind in dispersal, recruitment and survival rates, reproduction vegetatively versus
745 sexually, generation time, pollination biology, short and long-term impact of various
746 non-native invasive species, and short and long term habitat changes with differing
747 water levels. This information is required to understand population demographics,
748 reproductive ecology, dispersal and recruitment, and resiliency and will help inform best
749 management practices and a better informed definition of critical habitat. Research on
750 seed viability and germination requirements may be available from native plant/seed
751 suppliers in Manitoba. Finally, research is needed to determine whether the limited
752 number of populations and isolation of these populations represents a threat to genetic
753 diversity and/or population viability of the Canadian population as a whole. This type of
754 information will help determine the feasibility of restoration efforts and when/if
755 restoration efforts would be deemed necessary.

756
757 Additional inventory and monitoring work is a medium priority and required to know the
758 full extent (extent of occurrence and area of occupancy) of the population in Manitoba
759 and to obtain better estimates of population size. Regular monitoring will determine the
760 natural range of variability, trends, and health of the populations and will track whether
761 the population and distribution objectives are being met. Monitoring protocols should be
762 standardized and include metrics related to habitat quality, threats, plant health and
763 reproduction.

764
765

766 **7. Critical Habitat**

767
768 Critical habitat is defined in the *Species at Risk Act* (S.C.2002, c29) section 2(1) as “the
769 habitat that is necessary for the survival or recovery of a listed wildlife species and that
770 is identified as the species’ critical habitat in the recovery strategy or in an action plan
771 for the species”. Section 41 (1)(c) of *SARA* requires that recovery strategies include an
772 identification of the species’ critical habitat, to the extent possible, as well as examples
773 of activities that are likely to result in its destruction.

774
775 Critical habitat for Fascicled Ironweed is fully identified to the extent possible in this
776 recovery strategy, and is sufficient to achieve population and distribution objectives.
777 Additional critical habitat may be added or amended in the future if new or additional
778 information supports the inclusion or refinement of areas beyond those currently
779 identified (e.g., new sites become colonized, existing sites expand into adjacent areas,
780 historical populations are relocated, new information becomes available about habitat
781 requirements).

782 783 **7.1 Identification of the Species’ Critical Habitat**

784
785 Critical habitat for the Fascicled Ironweed is identified for the two known extant
786 populations based on best available information¹³ regarding the species’ location and
787 the associated biophysical attributes. Critical habitat is identified as all areas or types of
788 sites within critical habitat units (polygons; Appendix B) for the two known extant
789 populations that possess the following biophysical attributes (Table 4). Note that not all
790 biophysical attributes in Table 4 must be present in order for areas or types of sites to
791 be identified as critical habitat. If the area or type of site as described in Table 4 is
792 present and capable of supporting the species, the area or type of site is considered
793 critical habitat for the species.

794

¹³ Information on Fascicled Ironweed occurrences known to Environment and Climate Change Canada as of October 2019 was used in this recovery strategy.

795 **Table 4.** General summary of the areas and types of sites and biophysical attributes of critical
796 habitat for Fascicled Ironweed in Manitoba.

Area or type of site ^{a,b}	Biophysical attributes ^{b,c}
<ul style="list-style-type: none"> • open to semi-open native riparian areas including: <ul style="list-style-type: none"> • river terraces, oxbows, banks of creeks and wetlands; • moist to wet meadows; • floodplains, low-lying moist depressions or swales in tallgrass prairie • roadsides/ditches; • occasionally drier upland tallgrass prairie 	<ul style="list-style-type: none"> • tree species, when present, are deciduous (e.g. Green Ash, American Elm, Manitoba Maple, Eastern Cottonwood) • shrub layer is sparse to absent • herbaceous layer ranges from sparse to dense native vegetation <ul style="list-style-type: none"> • may have non-native species of plants amongst the native plants, • does not include thick dense vegetation cover from grasses and sedges or dense shade • soil is typically moist to wet clay but may tolerate other soil textures • habitat is seasonally flooded but may tolerate other levels of soil moisture and flooding

797 ^a Derived from Strother 2006, Foster and Reimer 2007, Manitoba Conservation unpubl. data 2019).

798 ^b Area or type of site: The area or type of site where the listed species naturally occurs or depends on in
799 order to carry out its life processes.

800 ^c Biophysical attributes: measurable properties or characteristics of the area or type of site. In essence,
801 biophysical attributes provide the greatest level of information about the area or type of site required to
802 support the life process requirements of the species.

803

804 7.1.1 Information and methods used to identify critical habitat

805

806 The approach used for identifying critical habitat for Fascicled Ironweed is based on a
807 decision tree developed by the Recovery Team for Plants at Risk in the Prairie
808 Provinces as guidance for identifying critical habitat for terrestrial and aquatic prairie
809 plant species at risk (see Appendix A in Environment Canada 2012 for the full decision
810 tree). Since Fascicled Ironweed is a perennial, reliably present, and easily detectable,
811 and occupies habitat that can be delineated in space and time, critical habitat was
812 identified as per criteria 3a of the decision tree. Accurate and precise occurrences that
813 had been confirmed in the last 25 years and habitat that had been confirmed in the past
814 five years were used. Occupied suitable habitat¹⁴ was mapped using the biophysical
815 attributes description, and included continuous or connected suitable habitat extending
816 away from the known occurrences. The river was also included in the mapping as it is
817 likely important for seed and propagule dispersal, maintaining connectivity and gene

¹⁴ Suitable habitat patches were delineated ex-situ using the best available satellite imagery (WorldView 02 imagery from May 2016 at 0.5 m spatial resolution at 1:3000 map scale, 10.2 m horizontal positional accuracy) following concepts of object-based segmentation (Jobin et al. 2008). Suitable habitat patches were visually identified based on vegetation community type using colour and texture. A 10 m buffer was placed around the delineated occupied habitat patches to account for the horizontal positional accuracy of the image (Paredes-Hernández et al. 2013, DigitalGlobe 2016) and temporal dynamics of habitat boundaries.

818 flow within the Rat River population, and allowing for natural expansion of the
819 population. For occurrences in roadside ditches where mapping boundaries was not
820 always obvious using the biophysical attributes description, the occupied suitable
821 habitat was bounded lengthwise by intersecting roads or road allowances/approaches
822 and widthwise by the road edge and the property line as this area contains uniform
823 conditions of hydrology and habitat (Environment Canada 2014). Critical habitat is
824 identified as all occupied suitable habitat and all natural biophysical attributes within a
825 300 m critical function zone extending from the outer boundary of the occupied suitable
826 habitat. Although the exact extent of habitat needed to surround Fascicled Ironweed
827 plants to fulfill the reproductive, dispersal and long-term survival needs of the population
828 is not fully documented, the 300 m critical function zone is based upon a detailed
829 literature review that examined edge-effects of various land use activities that could
830 affect resource availability and contribute to negative population growth for native prairie
831 plants generally (Henderson 2010b; Appendix B in Environment Canada 2012). In
832 addition, based on a detailed literature review that examined factors affecting the quality
833 of native prairie patches in the tallgrass prairie of Manitoba, the 300 m critical function
834 zone is appropriately applied to critical habitat identification of Fascicled Ironweed in
835 Manitoba (Environment and Climate Change Canada 2019 unpublished review). Thus,
836 to ensure the long-term persistence, and where feasible, the natural expansion, of
837 Fascicled Ironweed in Manitoba, the 300 m critical function zone is thought to be the
838 minimum distance needed to maintain the habitat required to meet the population and
839 distribution objectives. As new information on species' habitat requirements and
840 site-specific characteristics become available, this distance may be refined. The 300 m
841 critical function zone has been cropped so as to not include obvious existing human
842 developments and infrastructure, including cultivated lands, within the area identified as
843 critical habitat as these are not considered as critical habitat.

844

845 **7.1.2 Geographic information (Geographic Location of Areas Containing Critical** 846 **Habitat)**

847

848 The area containing critical habitat is approximately 1132 hectares (11.32 km²); no
849 critical habitat is on federal lands. Generalized geographic locations at the scale of
850 standardized 1x1 km grids and critical habitat unit polygons are provided in critical
851 habitat maps (Appendix B). All jurisdictions and landowners who are controlling surface
852 access to the area, or who are currently leasing and using parts of this area, may be
853 provided upon request with geo-referenced spatial data or large-format maps
854 delineating the boundaries of critical habitat displayed in Appendix B.

855

856 **7.2 Activities Likely to Result in the Destruction of Critical Habitat**

857

858 Destruction of critical habitat is assessed on a case by case basis. Destruction would
859 result if part of the critical habitat were degraded, either permanently or temporarily,
860 such that it would not serve its function when needed by the species. Destruction may
861 result from a single or multiple activities at one point in time (direct effect) or from the
862 cumulative effects of one or more activities over time (cumulative effect). Activities
863 described in Table 5 outline examples of activities likely to cause destruction of critical

864 habitat for Fascicled Ironweed; however, destructive activities are not limited to those
865 listed.

866
867

Table 5. Activities likely to result in the destruction of critical habitat.

Description of Activity	Description of Effect	Details of Effect
Compression or erosion of soils, which can be caused by activities such as: creation of trails and roads; motorized traffic; or concentration of livestock activity by the placement of bales, or establishment of new corrals or watering sites	Compression and erosion can damage soil structure and porosity, reduce water availability by increasing runoff and decreasing infiltration, prevent establishment of seedlings, or increase the likelihood of invasive non-native plants by disturbing native ground cover.	Related threats: 1.1 Housing & urban areas; 2.1 Annual & perennial non-timber crops; 2.3 Livestock farming & ranching; 4.1 Roads & railroads This activity must occur within the bounds of critical habitat to cause its destruction, can be a direct or cumulative effect, and is applicable at all times of the year, with the exception of winter months when the ground is snow covered and frozen solid (soil temperature below -10°C).
Covering of soils, which can be caused by activities such as: creation or expansion of permanent/ temporary structures such as land conversion to residential housing/developments; spreading of solid waste materials; or roadbed construction and certain road maintenance activities	Covering the soil prevents solar radiation and water infiltration needed for germination of seeds and survival of plants, such that critical habitat is destroyed.	Related threats: 1.1 Housing & urban areas; 2.1 Annual & Perennial Non-timber Crops; 2.3 Livestock Farming & Ranching; 4.1 Roads & railroads This activity must occur within the bounds of critical habitat to cause its destruction, is a direct effect, and is applicable at all times of the year.
Inversion/excavation/extraction of soils, which can be caused by activities such as: new or expanded cultivation; new/expansion of existing dugouts; certain road construction and maintenance activities or ditch deepening; residential development; pipeline installation; or removal of topsoil	Inverting, excavating or extracting soil results in the direct loss of critical habitat by removing or disturbing the substrate within which the plant grows, and altering the biophysical conditions (e.g. soil porosity, soil temperature, soil moisture) required for germination, establishment and growth of the Fascicled Ironweed. This activity can also lead to vegetation community change to one dominated by competitive invasive species.	Related threats: 1.1 Housing & urban areas; 2.1 Annual & Perennial Non-timber Crops; 2.3 Livestock Farming & Ranching; 4.1 Roads & railroads This activity must occur within the bounds of critical habitat to cause its destruction, can be both a direct and cumulative effect, and is applicable at all times of the year.
Alteration to hydrological regimes, which can be caused by activities such as: long-term or permanent inundation of upland habitat, or lack of seasonal flooding, or groundwater depletion, from	Fascicled Ironweed is adapted to moist soils and seasonal flooding events; changes to soil moisture or hydrology where the soil is too dry or too wet for an extended period of time can negatively affect the suitability of the habitat, affect plant growth and survival as well as seed	Related threats: 1.1 Housing & urban areas; 4.1 Roads & railroads; 7.2 Dams & water management/use This activity can occur within and outside the bounds of critical habitat to cause its

Description of Activity	Description of Effect	Details of Effect
impoundments downstream or alternately releases of water upstream; including but not limited to water or flow control structures (dams, ditches, dykes, diversions), drains, culvert installation, road widening or straightening; or residential developments	germination or seed bank viability, and change species composition in the area. Fascicled Ironweed grows in areas along the Rat River known to regularly flood seasonally in spring (sometimes in summer and fall) that have been inundated with water for periods of 3-6 weeks (Foster et al. 2007) and survived while other reports indicate it can tolerate a water depth of 18 inches decreasing over a period of four days (Shaw and Schmidt 2003); long-term or permanent flooding, however, of the habitat adjacent the river may be sufficient to alter that habitat enough to be unsuitable for survival or reestablishment. Alternately, altering hydrology to deplete groundwater or eliminate seasonal flooding long-term or permanently will likely result in conditions unsuitable for the species.	destruction, can be a direct or cumulative effect, and is applicable at all times of the year.
Indiscriminate application of fertilizers or pesticides	Herbicide and fertilizer can alter soil or water nutrient status, creating conditions suitable for some plant species and unsuitable for others, such that species composition in the surrounding plant community can change. Changes to soil or water nutrient status will also influence the outcome of interspecific competition for nutrients. Pesticide runoff and drift can alter plant and pollinator communities, thereby possibly reducing the capability of the habitat to support Fascicled Ironweed.	Related threats: 1.1 Housing & urban areas; 2.1 Annual & perennial non-timber crops; 4.1 Roads & railroads; 9.3 Agriculture & forestry effluents This activity can occur within and outside the bounds of critical habitat to cause its destruction (e.g. chemical drift, groundwater or overland flow of contaminated water), can be a direct or cumulative effect, and is applicable at all times of the year.
Deliberate introduction or promotion of invasive non-native plant species, which can be caused by activities such as: intentional dumping or spreading of feed bales containing viable seed of invasive non-native species; seeding invasive non-native species into critical habitat where the invasive non-native species did not already occur; use of motorized vehicles in critical habitat that are contaminated with invasive species material	Once established, invasive non-native plant species can alter hydrology, soil nutrient and moisture availability, and create dense shade or cover, resulting in direct competition with Fascicled Ironweed, such that population declines occur, effectively destroying the critical habitat. Critical habitat may be destroyed by invasive non-native species mentioned in Section 4.2 (threat 8.1), as well as by other noxious prohibited weeds and aggressive opportunistic species. It may also be destroyed by the following species which are not restricted by any legislation due to their economic value: Smooth Brome, Kentucky Bluegrass (<i>Poa pratensis</i>), Crested Wheatgrass (<i>Agropyron cristatum</i>), Yellow Sweet Clover (<i>Melilotus officinalis</i>), White Sweet Clover (<i>Melilotus alba</i>).	Related threats: 1.1 Housing & urban areas; 2.1 Annual & perennial non-timber crops; 2.3 Livestock farming & ranching; 4.1 Roads & railroads; 8.1 Invasive non-native species This activity can occur within or adjacent to the bounds of critical habitat to cause its destruction, can be a direct or a cumulative effect, and is applicable at all times of the year.

868

869 8. Measuring Progress

870

871 The performance indicators presented below provide a way to define and measure
872 progress toward achieving the population and distribution objectives. Beginning in 2020
873 and every five years thereafter, success of recovery strategy implementation will be
874 measured against the following performance indicators:

875

- 876 • The estimated population size and distribution of all extant populations and any
877 newly discovered or relocated populations are maintained, within the range of
878 natural variability.

879

880 9. Statement on Action Plans

881

882 One or more action plans will be posted on the Species at Risk Public Registry within
883 five years following the final posting of this recovery strategy.

884

885 10. References

- 886
887 Barnes, W.J. 1999. The rapid growth of a population of reed canarygrass (*Phalaris*
888 *arundinacea* L.) and its impact on some riverbottom herbs. *Journal of the Torrey*
889 *Botanical Society* 126: 133-138.
890
- 891 Brown, R.L. and J. Chenoweth. 2008. The effect of Glines Canyon Dam on
892 hydrochorous seed dispersal in the Elwha River. *Northwest Science* 82: 197-209.
893
- 894 Carthey, A.J.R, Fryirs, K.A., Ralph, T.J., Bu, H. and M.R. Leishman. 2016. How seed
895 traits predict floating times: a biophysical process model for hydrochorous seed
896 transport behaviour in fluvial systems. *Freshwater Biology* 61: 19-31.
897
- 898 Catford, J.A. and R. Jansson. 2014. Drowned, buried and carried away: effects of plant
899 traits on the distribution of native and alien species in riparian ecosystems. *New*
900 *Phytologist* 204: 19-36.
901
- 902 Channell, R. and M.V. Lomolino. 2000. Dynamic biogeography and conservation of
903 endangered species. *Nature* 403: 84–86.
904
- 905 COSEWIC. 2014. COSEWIC assessment and status report on the Fascicled Ironweed
906 *Vernonia fasciculata* in Canada. Committee on the Status of Endangered Wildlife in
907 Canada. Ottawa, ON. ix + 26 pp. Website: [https://wildlife-species.canada.ca/species-](https://wildlife-species.canada.ca/species-risk-registry/document/default_e.cfm?documentID=2758)
908 [risk-registry/document/default_e.cfm?documentID=2758](https://wildlife-species.canada.ca/species-risk-registry/document/default_e.cfm?documentID=2758) (accessed February 24, 2020).
909
- 910 COSEWIC. 2015. COSEWIC assessment process, categories and guidelines.
911 Committee on the Status of Endangered Wildlife in Canada. Website:
912 [https://www.canada.ca/content/dam/eccc/migration/cosewic-cosepac/94d0444d-369c-](https://www.canada.ca/content/dam/eccc/migration/cosewic-cosepac/94d0444d-369c-49ed-a586-ec00c3fef69b/assessment_process_and_criteria_e.pdf)
913 [49ed-a586-ec00c3fef69b/assessment_process_and_criteria_e.pdf](https://www.canada.ca/content/dam/eccc/migration/cosewic-cosepac/94d0444d-369c-49ed-a586-ec00c3fef69b/assessment_process_and_criteria_e.pdf)
914 (accessed February 24, 2020).
915
- 916 COSEWIC. 2020a. Definitions and abbreviations. Website:
917 <http://cosewic.ca/index.php/en-ca/about-us/definitions-abbreviations> (accessed
918 February 24, 2020).
919
- 920 COSEWIC. 2020b. Guidelines for use of the Index Area of Occupancy in COSEWIC
921 assessments. Website:
922 [http://cosewic.ca/index.php/en-ca/reports/preparing-status-reports/guidelines-index-](http://cosewic.ca/index.php/en-ca/reports/preparing-status-reports/guidelines-index-area-occupancy)
923 [area-occupancy](http://cosewic.ca/index.php/en-ca/reports/preparing-status-reports/guidelines-index-area-occupancy) (accessed February 24, 2020).
924
- 925 DigitalGlobe. 2016. Accuracy of WorldView Products: White Paper. Website:
926 [https://dgv4-cms-](https://dgv4-cms-production.s3.amazonaws.com/uploads/document/file/154/DG_ACCURACY_WP_V3.pdf)
927 [production.s3.amazonaws.com/uploads/document/file/154/DG_ACCURACY_WP_V3.p](https://dgv4-cms-production.s3.amazonaws.com/uploads/document/file/154/DG_ACCURACY_WP_V3.pdf)
928 [df](https://dgv4-cms-production.s3.amazonaws.com/uploads/document/file/154/DG_ACCURACY_WP_V3.pdf) (accessed June 28, 2019).
929

- 930 Douglas, M.R. and J.F. Tooker. 2015. Large-scale deployment of seed treatments has
931 driven rapid increase in use of neonicotinoid insecticides and preemptive pest
932 management in U.S. field crops. *Environmental Science and Technology*
933 49: 5088-5097.
- 934
935 Environment Canada. 2012. Recovery strategy for the Small-flowered Sand-verbena
936 (*Tripterocalyx micranthus*) in Canada. *Species at Risk Act Recovery Strategy Series*.
937 Environment Canada, Ottawa. v + 47 pp.
- 938
939 Environment Canada. 2014. Recovery strategy for the Small White Lady's-slipper
940 (*Cypripedium candidum*) in Canada. *Species at Risk Act Recovery Strategy Series*.
941 Environment Canada, Ottawa. v + 30 pp.
- 942
943 Foster, C. and C. Hamel. 2006. Rare species surveys of the Manitoba Conservation
944 Data Centre, 2005. MS Report 06-01. Manitoba Conservation Data Centre, Winnipeg,
945 Manitoba. 43 pp.
- 946
947 Foster, C. and E. Reimer. 2007. Rare plant surveys by the Manitoba Conservation Data
948 Centre, 2006. MS Report 07-01. Manitoba Conservation Data Centre, Winnipeg,
949 Manitoba. 53 pp. Website:
950 http://www.gov.mb.ca/conservation/cdc/pdf/field_report_2006.pdf (accessed
951 May 17 2019).
- 952
953 Friesen, C. and C. Murray. 2010. Rare Species Surveys and Stewardship Activities by
954 the Manitoba Conservation Data Centre, 2009. Report No. 2009-04. Manitoba
955 Conservation Data Centre, Winnipeg, Manitoba. 20 pp. Website:
956 http://www.gov.mb.ca/conservation/cdc/pdf/field_report_2009.pdf (accessed
957 May 17 2019).
- 958
959 Friesen, C. and C. Murray. 2011. Rare species surveys and stewardship activities by
960 the Manitoba Conservation Data Centre, 2010. Report No. 2010-01. Manitoba
961 Conservation Data Centre, Winnipeg, Manitoba. 24 pp. Website:
962 https://www.gov.mb.ca/sd/cdc/pdf/2010_final_report.pdf (accessed May 17 2019).
- 963
964 Gill, R.J. and N.E. Raine. 2014. Chronic impairment of bumblebee natural foraging
965 behavior induced by sublethal pesticide exposure. *Functional Ecology* 28: 1459-1471.
- 966
967 Godfray, H.C.J., Blacquiere, T., Field, L.M., Hails, R.S., Petrokofsky, G., Potts, S.G.,
968 Raine, N.E., Vanbergen, A.J. and A.R. McLean. 2014. A restatement of the natural
969 science evidence base concerning neonicotinoid insecticides and insect pollinators.
970 *Proceedings of The Royal Society Biological Sciences* 281: 20140558.
- 971
972 Great Plains Flora Association. 1986. *Flora of the Great Plains*. University Press of
973 Kansas, Kansas. 1402 pp.
- 974

- 975 Groves, J. 2010. Fluvial seed dispersal processes and the sustainability of riparian
976 vegetation. PhD thesis. University of Canberra, Canberra, ACT.
977
- 978 Henderson, D.C. 2010a. Occupancy survey guidelines for Prairie Plant species at risk.
979 Environment Canada, Prairie and Northern Region, Canadian Wildlife Service.
980 Edmonton, AB. Website:
981 www.npss.sk.ca/docs/2_pdf/Rare_Plant_Occupancy_Survey_Guidelines.pdf (accessed
982 July 18, 2019).
983
- 984 Henderson, D.C. 2010b. Set-back distance and timing restriction guidelines for prairie
985 plant species at risk. Environment Canada, Prairie and Northern Region, Canadian
986 Wildlife Service. Edmonton AB. Website:
987 https://www.npss.sk.ca/rsu_docs/documents/plant-setback-guidelines-nov-15-2010.pdf
988 (accessed July 18, 2019).
989
- 990 Hubbard, D.E. and A. Boe. 1988. IVDDM and chemical constituents in selected species
991 of wetland plants. Proceedings South Dakota Academy of Science 67: 44-58.
992
- 993 Jansson, R., Zinko, U., Merritt, D.M. and C. Nilsson. 2005. Hydrochory increases
994 riparian plant species richness: a comparison between a free-flowing and a regulated
995 river. Journal of Ecology 93: 1094-1103.
996
- 997 Jobin, B., Labrecque, S., Grenier, M. and G. Falardeau. 2008. Object-based
998 classification as an alternative approach to the traditional pixel-based classification to
999 identify potential habitat of the grasshopper sparrow. Environmental Management
1000 41: 20-31.
1001
- 1002 Jones, S.B. 1972. A systematic study of the Fasciculatae group of Vernonia
1003 (Compositae). Brittonia 24: 28-45.
1004
- 1005 Kartesz, J.T. 2015. *North America Plant Atlas*. Chapel Hill, N.C. [maps generated from
1006 Kartesz, J.T. 2015. Floristic Synthesis of North America, Version 1.0. Biota of North
1007 America Program (BONAP). (in press)]. Website: <http://www.bonap.org/napa.html>
1008 (accessed May 22, 2019).
1009
- 1010 Kearnes, C.A., Inouye, D.W. and N.M. Waser. 1998. Endangered mutualisms: The
1011 conservation of plant-pollinator interactions. Annual Review of Ecology and Systematics
1012 29: 83–112.
1013
- 1014 Lavergne, S. and J. Molofsky. 2007. Increased genetic variation and evolutionary
1015 potential drive the success of an invasive grass. Proceedings of the National Academy
1016 of Sciences 104: 3883-3888.
1017
- 1018 Lesica, P., Moore, G., Peterson, K.M. and J.H. Rumely. 1984. Vascular Plants of
1019 Limited Distribution in Montana. Montana Academy of Sciences, Supplement to the
1020 Proceedings, Vol. 43, Monograph 2.
1021

- 1022 Medley, M. 1993. An Annotated Catalog of the Known and Reported Vascular Flora of
1023 Kentucky. Ph.D. Dissertation, Univ. of Louisville, Louisville.
1024
- 1025 Merritt, D.M. and E.E. Wohl. 2002. Processes governing hydrochory along rivers:
1026 hydraulics, hydrology, and dispersal phenology. *Ecological Applications* 12: 1071-1087.
1027
- 1028 Merritt, D.M. and E.E. Wohl. 2006. Plant dispersal along rivers fragmented by dams.
1029 *River Research and Applications* 22: 1-26.
1030
- 1031 Montana Natural Heritage Program. 2019. Montana Field Guide: Fascicled Ironweed –
1032 *Vernonia fasciculata* ssp. *corymbosa*. Website:
1033 <http://fieldguide.mt.gov/speciesDetail.aspx?elcode=PDAST9S091> (accessed
1034 May 17 2019).
1035
- 1036 Murray, C. 2019. Personal Communications. Information Manager, Manitoba
1037 Conservation Data Centre, Manitoba Conservation.
1038
- 1039 Murray, C. and C. Friesen. 2012. Manitoba Conservation Data Centre Surveys and
1040 Stewardship Activities, 2011. Report No. 2012-01. Manitoba Conservation Data Centre,
1041 Winnipeg, Manitoba. 24 pp. Website:
1042 https://www.gov.mb.ca/sd/cdc/pdf/2012_field_report.pdf (accessed May 17 2019)
1043
- 1044 Murray, C. 2013. Manitoba Conservation Data Centre Surveys and Stewardship
1045 Activities, 2012. Report No. 2013-01. Manitoba Conservation Data Centre, Winnipeg,
1046 Manitoba. 30 pp. Website:
1047 https://www.gov.mb.ca/sd/cdc/pdf/2012_field_report_2013_01.pdf
1048 (accessed May 17 2019)
1049
- 1050 Murray, C. 2014. Manitoba Conservation Data Centre Surveys and Stewardship
1051 Activities, 2013. Report No. 2014-01. Manitoba Conservation Data Centre, Winnipeg,
1052 Manitoba. v+41 pp. Website: https://www.gov.mb.ca/sd/cdc/pdf/2014_field_report.pdf
1053 (accessed May 17 2019).
1054
- 1055 Murray, C. and C. Church 2015. Manitoba Conservation Data Centre surveys and
1056 stewardship activities, 2014. Report No. 2015-01. Manitoba Conservation Data Centre,
1057 Winnipeg, Manitoba. v+47 pp. Website:
1058 https://www.gov.mb.ca/sd/cdc/pdf/manitoba_conservation_data_centre_2014.pdf
1059 (accessed May 17 2019).
1060
- 1061 Murray, C. and C. Church. 2017. Manitoba Conservation Data Centre surveys and
1062 stewardship activities, 2015. Report No. 2017-02. Manitoba Conservation Data Centre,
1063 Winnipeg, Manitoba. lii + 55 pp.
1064
- 1065 Nabhan, G.P. and S.L. Buchmann. 1997. Services provided by pollinators. In: *Nature's*
1066 *Services Societal Dependence on Natural Ecosystems*. G. C. Daily. Island Press.
1067 412 pp.

- 1068
1069 NatureServe. 2020a. Habitat-based Plant Element Occurrence Delimitation Guidance,
1070 1 October 2004. Version 7.1. NatureServe, Arlington, Virginia. Website:
1071 http://explorer.natureserve.org/decision_tree.htm (accessed January 13, 2020).
1072
- 1073 NatureServe. 2020b. NatureServe Explorer: An online encyclopedia of life [web
1074 application]. Version 7.1. NatureServe, Arlington, Virginia. Website:
1075 <http://explorer.natureserve.org/> (accessed January 13, 2020).
1076
- 1077 NatureServe. 2020c. Ranking species occurrences – A generic approach,
1078 11 January 2008. Version 7.1. NatureServe, Arlington, Virginia. Website:
1079 <http://explorer.natureserve.org/eorankguide.htm> (accessed January 13, 2020).
1080
- 1081 Paredes-Hernández, C.U., Salinas-Castillo, W.E., Guevara-Cortina, F. and X.
1082 Martínez-Becerra. 2013. Horizontal positional accuracy of Google Earth’s imagery
1083 over rural areas: a study case in Tamaulipas, Mexico. *Boletim de Ciências*
1084 *Geodésicas* 19: 5880601.
1085
- 1086 Scheper, J., Reemer, M., Kats, R., Ozinga, W.A., Linden, G.T.J., Schaminee, J.H.J.,
1087 Siepel, H. and D. Kleijn. 2014. Museum specimens reveal loss of pollen host plants as
1088 key factor driving wild bee decline in The Netherlands. *Proceedings of the National*
1089 *Academy of Sciences* 111: 17552-17557.
1090
- 1091 Shaw, D. and R. Schmidt. 2003. *Plants for stormwater design*. Minnesota Pollution
1092 Control Agency, Saint Paul, Minnesota.
1093
- 1094 Sluis, W.J. and J. Tandarich. 2004. Siltation and hydrologic regime determine species
1095 composition in herbaceous floodplain communities. *Plant Ecology* 173: 115-124.
1096
- 1097 Smith, E.B. 1973. An annotated list of the Compositae of Arkansas. *Castanea*
1098 38: 79-109.
1099
- 1100 Smith, E.B. 1988. *An atlas and annotated list of the vascular plants of Arkansas*, 2d ed.
1101 University of Arkansas, Fayetteville. 592 pp.
1102
- 1103 Strother, J.L. 2006. *Vernonia*. In: *Flora of North America* Editorial Committee, Eds.
1104 1993+. *Flora of North America North of Mexico*. 16+ vols. New York and Oxford. Vol.
1105 19. Pp. 206-213.
1106
- 1107 Uowolo, A.L., Binkley, D. and E. Carol Adair. 2005. Plant diversity in riparian forests in
1108 northwest Colorado: effects of time and river regulation. *Forest Ecology and*
1109 *Management* 218: 107-114.
1110
- 1111 USDA. 2019. The PLANTS Database. National Plant Data Center, Baton Rouge, LA.
1112 Website: <http://plants.usda.gov> (accessed July 18, 2019).
1113

- 1114 Vucetich, J.A. and T.A.Waite. 2003. Spatial patterns of demography and genetic
1115 processes across the species' range: Null hypothesis for landscape conservation
1116 genetics. *Conservation Genetics* 4: 639-645.
1117
- 1118 White, M. 2012. *Vernonia fasciculata* (Asteraceae): New to the flora of Texas.
1119 *Phytoneuron* 2012-90: 1-6. Website: [http://www.phytoneuron.net/90PhytoN-](http://www.phytoneuron.net/90PhytoN-VernoniaNew.pdf)
1120 [VernoniaNew.pdf](http://www.phytoneuron.net/90PhytoN-VernoniaNew.pdf) (accessed July 18, 2019).
1121
1122

1123 **Appendix A: Summary of Fascicled Ironweed Populations in Canada**

1124


1125 **Table A1.** Summary of Fascicled Ironweed Populations in Canada. Light grey shading indicates the population is extirpated or
 1126 historical.

Population Name [EO_ID]	First Observed	Last Observed	Recent Survey Estimate [Year]	Highest Estimate [Year]	Status	Threats	Notes
MANITOBA							
Morris [1750]	1896	1953	0 [2013]	1 clump [1953]	Historical but believed extirpated	N/A	Surveys in the area of Morris in 2005, 2006, 2010, and 2013 found no plants (MB Conservation Data Centre, unpubl. data 2019). Habitat description from 1896 was from along what is now the Morris River and another location from 1953 was "grassy road ditch"; COSEWIC has this population as extirpated and states it was eliminated by the conversion of native pasture to cropped field (Foster and Reimer 2007, Friesen and Murray 2011, COSEWIC 2014). A 12 km portion of the Morris River has also been searched but no Fascicled Ironweed found.
Mile Road 4W [4867]	1995	2014	~11 plants containing ~83 stems [2014]	35 plants plus 27 stems [2010]	Current	4.1; 9.3	The upland prairie portion of this population was extirpated between 1995-2006 through cultivation; plants are now restricted to a municipal ditch and drain with the ditch being cleared of vegetation sometime between 2010 and 2013 (Foster and Reimer 2007, Friesen and Murray 2011, COSEWIC 2014, MB Conservation Data Centre, unpublished data 2019).

Population Name [EO_ID]	First Observed	Last Observed	Recent Survey Estimate [Year]	Highest Estimate [Year]	Status	Threats	Notes
Rat River [2709]	1950	2019	>1,766 plants plus >15,911 stems [2018]; additional >2066 plants and >6361 stems [2019]	>1,766 plants plus >15,911 stems [2018]; additional >2066 plants and >6361 stems [2019]	Current	1.1; 2.1; 2.3; 4.1; 7.2; 8.1; 9.3	Population estimates between years are not comparable as different portions of the population were surveyed each year. The most recent survey effort in 2018 and 2019 does not include the entire Rat River population. Some portions have been revisited multiple times but the majority have not. Note that Provincial Road 200 (PR 200) was considered separately in the COSEWIC (2014) report but is now recognized as part of Rat River (2709).
SASKATCHEWAN							
Weyburn [16246]	pre-1949	pre-1949	0 [2017]	>1 (pre-1949)	Historical	Unknown	Population has never been relocated; original herbarium record is very vague with respect to the location (ie. "Weyburn prairie"). Targeted searches have occurred in seemingly suitable habitat in the Weyburn area with no success.


1127
1128

Appendix B: Critical Habitat for Fascicled Ironweed in Canada



1129
1130
1131
1132
1133

Figure B1. Critical habitat for Fascicled Ironweed in Manitoba (Mile Road 4W [EO 4867] population as described in Table A1) is represented by the yellow shaded units, where the criteria set out in Section 7.1 are met. The 1 km x 1 km UTM grid overlay shown on this figure is a standardized national grid system that indicates the general geographic area containing critical habitat. Areas outside of the yellow shaded units do not contain critical habitat.



1134
1135
1136
1137
1138

Figure B2. Critical habitat for Fascicled Ironweed in Manitoba (Rat River [EO 2709] population as described in Table A1) is represented by the yellow shaded units, where the criteria set out in Section 7.1 are met. The 1 km x 1 km UTM grid overlay shown on this figure is a standardized national grid system that indicates the general geographic area containing critical habitat. Areas outside of the yellow shaded units do not contain critical habitat.

1139 **Appendix C: Effects on the Environment and Other Species**

1140

1141 A strategic environmental assessment (SEA) is conducted on all SARA recovery
1142 planning documents, in accordance with the [Cabinet Directive on the Environmental
1143 Assessment of Policy, Plan and Program Proposals](#)¹⁵. The purpose of a SEA is to
1144 incorporate environmental considerations into the development of public policies, plans,
1145 and program proposals to support environmentally sound decision-making and to
1146 evaluate whether the outcomes of a recovery planning document could affect any
1147 component of the environment or any of the [Federal Sustainable Development
1148 Strategy](#)'s¹⁶ (FSDS) goals and targets.

1149

1150 Recovery planning is intended to benefit species at risk and biodiversity in general.
1151 However, it is recognized that strategies may also inadvertently lead to environmental
1152 effects beyond the intended benefits. The planning process based on national
1153 guidelines directly incorporates consideration of all environmental effects, with a
1154 particular focus on possible impacts upon non-target species or habitats. The results of
1155 the SEA are incorporated directly into the strategy itself, but are also summarized below
1156 in this statement.

1157

1158 The potential for the strategy to inadvertently lead to adverse effects on other species
1159 was considered, including the many federal species at risk and provincially rare species
1160 that are found in habitat occupied by Fascicled Ironweed (Table C1). Broad approaches
1161 to recovery for the Fascicled Ironweed include communication/ collaboration
1162 /engagement, habitat assessment/ management/ stewardship activities, inventory/
1163 monitoring, and research. These activities are aimed at maintaining or improving
1164 riparian and wet meadow habitats. For the most part, managing for healthy native
1165 ecosystems, maintaining or improving native habitat, and conservation or stewardship
1166 actions will benefit non-target species, natural communities, and ecological processes.
1167 As a general rule, management actions that incorporate or mimic natural processes are
1168 natural components of prairie ecosystems and are not likely to negatively impact the
1169 persistence of other native species particularly if the timing, intensity and frequency
1170 mimic those natural processes (Samson and Knopf 1994). However, some
1171 management practices, and some forms of integrated weed management, have the
1172 potential to affect some species negatively in the short or long-term. Therefore, it is
1173 important that management actions resulting from recovery strategies, action plans and
1174 beneficial management plans are developed with experts from an ecosystem
1175 perspective (including development of multi-species action plans and ecosystem
1176 beneficial management plans), incorporating as many species' needs as possible, and
1177 evaluating the ecological risks of any action, in order to reduce any possible negative
1178 effects to other species; in some cases, this may need to be done on a site-specific
1179 basis. An ongoing monitoring program should also be in place to evaluate baseline
1180 conditions, and the short and long term effects of management actions on the
1181 ecosystem and individual species at risk so efforts can be adapted if negative impacts

¹⁵ www.canada.ca/en/environmental-assessment-agency/programs/strategic-environmental-assessment/cabinet-directive-environmental-assessment-policy-plan-program-proposals.html

¹⁶ www.fsds-sfdd.ca/index.html#/en/goals/

1182 are observed. Efforts should be coordinated with recovery teams and organizations
 1183 working in the tallgrass prairie ecosystem to ensure the most efficient use of resources
 1184 and to prevent duplication of effort or conflicts with research.

1185
 1186 The SEA concluded that this strategy will not entail significant adverse environmental
 1187 effects. For further details see the following sections: 3.3 Needs of the Fascicled
 1188 Ironweed, 4. Threats, and 6. Broad strategies and general approaches to meet
 1189 objectives.

1190
 1191 **Table C1.** Federal species at risk that co-occur, or may co-occur, in areas occupied by
 1192 Fascicled Ironweed, as identified by Conservation Data Centre records within a 1 km radius of
 1193 Fascicled Ironweed.

Common Name	Scientific name	SARA status	COSEWIC status
Barn Swallow	<i>Hirundo rustica</i>	Threatened	Threatened
Bobolink	<i>Dolichonyx oryzivorus</i>	Threatened	Threatened
Chimney Swift	<i>Chaetura pelagica</i>	Threatened	Threatened
Eastern Wood-pewee	<i>Contopus virens</i>	Special Concern	Special Concern
Golden-winged Warbler	<i>Vermivora chrysoptera</i>	Threatened	Threatened
Mapleleaf Mussel	<i>Quadrula quadrula</i>	Endangered	Threatened
Monarch	<i>Danaus plexippus</i>	Special Concern	Endangered
Northern Leopard Frog	<i>Lithobates pipiens</i>	Special Concern	Special Concern
Snapping Turtle	<i>Chelydra serpentina</i>	Special Concern	Special Concern

1194